

**Darwin Initiative Award 15/036: Monitoring and Managing Biodiversity Loss in South-east Africa's Montane Ecosystems**

# **THE BIODIVERSITY AND CONSERVATION OF MOUNT CHIPERONE, MOZAMBIQUE**



July 2007

Jonathan Timberlake, Julian Bayliss, Tereza Alves, Susana Baena, Jorge Francisco, Tim Harris, Camila da Sousa



Forestry Research  
Institute of Malawi



## LIST OF CONTENTS

1.	INTRODUCTION	3
2.	STUDY AREA	3
	Geology and Geomorphology	5
	Climate	5
	Landuse	5
3.	PREVIOUS STUDIES	6
4.	VEGETATION TYPES	6
	Vegetation Mapping	7
	Historical Change	11
	Vegetation Types	11
	Summit Thicket	12
	High-Altitude Forest	12
	Medium-Altitude Forest	13
	Miombo and Similar Woodland Types	13
	Disturbed Woodland and Fallows	14
	Rocky Outcrops	15
5.	BOTANY	15
6.	ZOOLOGY	16
	Local Hunter Records	17
	Small Mammals	18
	Birds	18
	Reptiles and Amphibians	20
	Lepidoptera	20
	Coleoptera and Hemiptera	22
7.	THREATS AND CONSERVATION ISSUES	23
8.	RECOMMENDATIONS	24
	Research Issues	24
9.	CONCLUSIONS	24
10.	REFERENCES	25
	ANNEX 1. Participants on Mt Chipirone Expedition, Nov/Dec 2006	27
	ANNEX 2. Plant checklist from Mt Chipirone above 800 m	28
	ANNEX 3. List of birds seen on Mt Chipirone, 15–18 December 2005	32

## 1. INTRODUCTION

A scientific expedition to Mount Chiperone in northern Mozambique was carried out from 22 November to 5 December 2006. The expedition was funded under a Darwin Initiative grant to the Royal Botanic Gardens, Kew – "Monitoring and Managing Biodiversity Loss on South-East Africa's Montane Ecosystems". The expedition was a collaborative effort between Kew, the Instituto de Investigação Agrária de Moçambique (IIAM), the Mulanje Mountain Conservation Trust (MMCT), and the Forest Research Institute of Malawi (FRIM). A full list of participants is given in Appendix 1.

The objectives of the expedition and study were:

1. To undertake botanical and vegetation field survey of Mount Chiperone
2. To gather additional zoological information on the mountain
3. To train a team of Mozambican and Malawian biologists in botanical and vegetation survey techniques
4. To assess the extent and status and threats to the moist forest and other biodiversity on the mountain
5. Based on gathered field data, to develop species and habitat recovery plans.

This report attempts to document what is known on the biodiversity and physical attributes of Mount Chiperone, to present the results of the expedition, and to outline the threats to that biodiversity. It also outlines conservation management issues, with particular reference to moist forest above 1000 m altitude, and gives some conservation recommendations. Species and habitat recovery plans will be outlined in another report towards the end of the project.

## 2. STUDY AREA

### **Location**

Mount Chiperone in northern Mozambique is a semi-isolated peak situated some 50 km south of the Mount Mulanje massif in southern Malawi. It lies in Milange District of Zambézia Province, 40 km SSW of the District Centre of Milange. The area of Mount Chiperone above 600 m covers about 100 km<sup>2</sup>, with around 4915 ha (in planimetric view) above 800 m altitude, 2770 ha above 1000 m, and only 110 ha over 1800 m. The massif is centred on 16°29'S, 35°43'E, with the highest point at 2054 m (c. 16°28'44"S, 35°42'88"E).

The surrounding plains to the north form part of the central African plateau, here lying at around 400–450 m altitude, while the land to the south falls away rapidly to 200–350 m towards the coast some 200 km away from the Zambezi Delta. The Shire valley, part of the East African Rift Valley, lies 35 km to the west at an altitude at this point of 50–100 m.

Starting from a road-head in Sabelua village (16°30'48"S, 35°46'08"E, alt. 400 m) on the SE slopes, expedition members walked up the south-eastern flank through cleared and regenerating woodland vegetation, across two ridges, and established a base camp just inside the nearest patch of moist forest (16°30'35.4"S, 35°43'42.7"E) at an altitude of 1029 m. Most of the survey work was concentrated in the forest and miombo woodland within relatively easy access of the camp. In addition, reconnaissance trips were made up along the NW–SE trending summit ridge and on the eastern slopes above Marega village (16°28'1.4"S, 35°45'45.1"E, alt. 502 m). Collecting localities and areas visited are shown in Figure 1.




Figure 1. Map of Chipirone area and collecting localities.

Logistics and the full itinerary are given in a separate report (J. Bayliss, Trip Report – Mount Chipirone Expedition, 22 Nov–5 Dec 2006).

### **Geology and Geomorphology**

Mount Chipirone, a semi-isolated peak, appears to rise rapidly out of the surrounding plateau, and has a distinctive small pointed peak when viewed from the north or east. There is a NW–SE trending ridge, with a main peak (2054 m, ONC charts) slightly towards the NW end. However published figures vary slightly, from 2017 m [Google Earth] to 2065 m [1:250,000 map]. Other minor high points are seen along the ridge, some of which are almost devoid of woody cover. Otherwise the slopes coming off the ridge are steep and mostly covered in closed moist forest.

The mountain is composed of fairly recent syenite (Jurassic/Cretaceous period, c.150 Mya), intruded the surrounding country rock, which comprises migmatites (metamorphic rock injected with igneous material) of the Namarroi series (850–1100 Mya). In this respect, Chipirone differs from most of the other massifs or hills in northern Mozambique, which are composed of migmatites or granites (e.g. Mt Namuli, the inselbergs around Ribaué and Nampula). The exceptions are Mt Tembe at Morrumbala and Mount Tumbine above Milange town, and much further to the south, a large part of Mount Gorongosa. Mt Mulanje, formed 130 Mya at approximately the same time as Chipirone, is composed of syenite, quartz-syenite and granite (Garson & Walshaw 1969, Eastwood 1988), nomenclature depending on the level of quartz (syenite if quartz is essentially absent).

The main drainage off the mountain is to the south west. These small rivers, the Rio Macololo (becoming the Rio Metambe) and the Rio Muse, are probably not perennial. They both drain into the Shire River in southern Malawi above the town of Nsanje.

### **Climate**

Climatic data on the area are not available. From coarse-scale maps, mean annual rainfall is around 1400 mm/year. Meteorological data from Milange, 40 km away (Kassam et al. 1981), give a mean annual rainfall of 1733.9 mm (28 years), with a mean annual temperature of 23°C, mean maximum of 28.9°C and mean minimum of 17.1°C. Vegetation on the lower slopes of the mountain, however, suggests that rainfall is somewhat lower, perhaps around 1000 mm/year.

### **Landuse**

There are two villages and settlements at the base of Mt Chipirone on the south and south-eastern slopes, Sabelua and Marega. The peneplain here has been extensively cleared and is cultivated at subsistence level. The northern, north-eastern and western slopes seem little affected by human activity, although they were not visited on the ground. On the lower slopes on the south and south-east there has been patchy clearance for cultivation of maize, cassava and beans, and extensive burning, particularly in the Marega area. The landscape of Mt. Chipirone is a mosaic of cultivation, disturbed woodland and fallows/regenerating woodland of various ages. This continues through the miombo woodland belt until the edge of the moist forest. There is virtually no cultivation or cleared patches above 1000 m altitude on the southern and south-western slopes, but cleared patches are found up to 1400 m on the (presumably moister or more mesic) south-eastern slopes.

The moist forest itself appears very little disturbed, apart from fire along the boundaries, particularly marked in the gullies. This lack of anthropogenic disturbance is possibly due to the local belief of malevolent spirits residing in the forest.

Water was identified as a particular problem for villages at the base of the mountain. There are a few small streams running off, but during the dry season these dry up or become a trickle. There are apparently no wells or boreholes in the vicinity.

### 3. PREVIOUS STUDIES

It appears that very little published information is available on Mt Chipero and the immediate surrounding area. This is rather surprising considering the wealth of information – geological, biological, historical – on Mt Mulanje in Malawi, just 50 km away. However, rarely did researchers and others based in Malawi extend their study or visit across the border, even though in the first part of the 20th century access to this part of Malawi was often by train through Mozambique and up the Shire Valley. Dixey, the Nyasaland Government Geologist, for example, does not mention Chipero in his study on the "Mlanje Mountains of Nyasaland" (Dixey 1927), and Jim Chapman, a Forester who spent many years working and collecting on and around Mt Mulanje, has also not visited (pers. comm. 2007). Vincent, in his mammoth bird-collecting trip across southern Malawi and northern Mozambique, did not visit, but saw the "island" Mt Chipero from afar and said it "..... should hold much interest to the naturalist". The main knowledge of Chipero among many residents of Malawi would appear to be through the so-called "chipero" weather, a 5-day misty weather system that is reputed to generate over Chipero and which then moves towards Mulanje, bringing light rainfall. Such weather is of particular significance and benefit to the tea plantations there.

The first recorded biological collecting from Mount Chipero was of birds, carried out from 25–31 July 1950 by Jali Makawa, a collector for the famous ornithologist C.W. Benson (reported in Benson 1950 and Spottiswoode et al. 2006). Over 6 days at an altitude of around 1500 m Makawa collected at least one specimen of 18 species, with 9 additional sight records, including a number of threatened or range-restricted species (see Parker 2001). Eight were new records for Mozambique at that time. He reported that there was extensive evergreen forest on the eastern slopes (3 square miles or 780 ha), more extensive than anything in southern Malawi. There are reports of small mammals and birds being collected around 2002 by persons from the Chicago Field Museum and Department of Biology of Universidade Eduardo Mondlane in Maputo, but results are not yet available. In December 2005, Claire Spottiswoode, Hassam Patel, Eric Herrmann and Julian Bayliss recorded forest birds in the same part of the mountain visited by the present expedition, and also collected some of the main plants. A report on this study and an assessment of forest extent is given in Spottiswoode et al. (2007).

### 4. VEGETATION TYPES

At a regional scale the Chipero massif is relatively small, but even so both Wild and Barbosa (1967) in the vegetation map of the Flora Zambesiaca area, and Frank White (1983) in his major work on African vegetation, depict it. White shows Chipero as a forest patch of East African coastal mosaic (type 16b), similar to that on other montane massifs in northern Mozambique, and surrounded by Drier Zambezian Miombo Woodland (type 26). The more detailed map by Wild & Barbosa (1967), which formed the basis of White's map, shows it as Moist Evergreen Forest (medium and low altitude), surrounded by *Brachystegia spiciformis*–*Julbernardia* Woodland. Our observations support this, although little *B. spiciformis* was seen in the surrounding woodlands, and the moist forest was primarily medium altitude, not low altitude.

Earlier studies include Barbosa's (1952) study on the vegetation of Zambézia Province. He describes the vegetation of Mt Chipero, along with that on other massifs such as Mt Mabu and

Morrumbala, as Unit 1, Moist Tropical Montane Forest (rain and clouds). He says that in general trees are evergreen, 18–20 m high with 3 or 4 strata, and forest is only found at over 1200 m altitude. The herbaceous layer is poor, but ferns are common. He also points out that additional moisture is available to these forests through clouds being formed as the prevailing moist south-easterly airflow is forced over the mountains and cools. Above a certain, unspecified, altitude temperatures are sufficiently cool that a xerophytic thicket vegetation is encountered dominated by species from the Ericaceae and Proteaceae families. This is what the present survey found at around 1900–2000 m. According to Barbosa, typical moist forest species include: *Albizia gummifera*, *Anthocleista grandiflora*, *Coffea ligustroides*, *Cussonia arborea*, *Entada rheedei*, *Harungana madagascariensis*, *Heteropyxis natalensis*, *Macaranga* spp., *Maesa lanceolata*, *Newtonia buchananii*, *Oxyanthus speciosus*, *Parinari curatellifolia*, *Parinari excelsa*, *Smilax anceps*, *Trichila dregeana*, *Vitex* spp., and the herbs *Afromomum* spp., *Costus* sp, *Ensete* sp. and *Piper umbellatum*. A number of these species, although not all, were noted in the forests of Mt Chiperone.

Pedro & Barbosa (1955) produced a map of the vegetation of Mozambique, which later formed the basis of the Mozambique section of Wild & Barbosa's Flora Zambesiaca map. They do not give details of vegetation in our study area as apparently this was only seen from afar, but they state that vegetation at 1000–1800 m is part of Complex 79 (Montane zones of Zambézia–Niassa), while those parts above 1800 m fall into Complex 80 (Subalpine zones of Zambézia).

### Vegetation Mapping

Vegetation mapping, specifically determination of the extent of moist forest, was carried out using two separate techniques. These were manual interpretation of remotely-sensed data supported by study of available air photos, and the supervised classification of a combination of Landsat ETM+ and ASTER digital imagery. In addition, the historical air photos were used to determine approximate moist forest extent in 1969. These studies are elaborated upon below.

Forest is here defined as a continuous stand of trees with interlocking crowns, mostly over 10 m in height. It differs from the FAO definition, which covers most stands of woody plants including what we would term woodland.

The only air photos available were from 1969 at a scale of 1:46,500. Unfortunately cloud cover over the peak and shadows from it reduced their usefulness. Orientation once up the mountain was difficult owing to the steeply-dissected terrain, lack of locatable landmarks, and uncertainty on extent of vegetation clearance and infrastructural changes since 1969.

Air photos of Mt Chiperone, 1:46,500 scale, 1969

path 76	39/ 168–171
path 77	39/ 193–195

#### a) *Manual Interpretation*


Manual interpretation of forest extent was carried out using a grey-scale Landsat ETM image from May 2002 as the base, and supported by use of 1969 air photos and field knowledge from the southern slopes. This polygon was digitised and put into a GIS (Figure 3). Based on this, forest extent was calculated to be 1717 ha (Table 1).

#### b) *Digital Classification*

Initially a draft vegetation map was made based on an unsupervised classification (maximum likelihood algorithm, applied to a 6-band stack image) of a Landsat ETM+ image acquired from May 2002 (path 167, row 071). Nine classes were recognised, including two different forest types

with an additional 'shadow' class. Based on this initial interpretation, it was calculated that 2063 ha of forest was present, with approximately 92% of this lying above 800 m altitude.

Following fieldwork and the recording of 170 ground control points (GPS readings in areas clearly either forest, woodland or cleared), a final vegetation map was developed using a Landsat ETM+ image acquired from May 2002 (30 m resolution, Figure 2a) and an ASTER image acquired in September 2001 (15 m resolution, Figure 2b). Both images (path/row 167/071) were registered to UTM Zone 36 S (WGS 84) with radiometric and geometric correction. Following an initial inspection of both images, Landsat proved to discriminate different vegetation types more accurately despite its coarser resolution. In order to gain information from the ASTER image, the NDVI was developed and stacked to the 6-band Landsat product. A supervised classification was then performed on the combination of original Landsat bands and NDVI. The following vegetation types were separated: forest, open forest, woodland, open savanna and cultivation. High-altitude forest was not spectrally different from medium-altitude forest, therefore a threshold of 1600 m was used to differentiate them. The resulting map is shown in Figure 4, with the extent of forest types given in Table 1.



Figures 2a and 2b. Mt Chipirone area – Landsat ETM+ image, May 2002 (left) and ASTER satellite image, September 2001 (right).

Relatively minor differences can be seen between the extent of forest as determined using the two techniques. As forest was hardly noted in the field below 1000 m, at least on the southern and eastern slopes, it is possible that the forest category also includes some miombo (*Brachystegia*-dominant) woodland areas.

Table 1. Forest extent (2002) on Mt Chipirone determined using different methods.

	Forest extent (ha)	
	Supervised	Manual
Open forest - medium altitude	241.6	–
Open forest - high altitude	165.4	–
Forest - medium altitude	1051.5	1307
Forest - high altitude	176.6	410
TOTAL	1635.1	1717





## VEGETATION MAP - CHIPERONE






Figure 3. Extent of forest on Mt Chipirone (visual interpretation).



# VEGETATION MAP - CHIPERONE



**LEGEND**

**Vegetation types**

- Cultivation
- Forest
- Montane forest
- Open forest
- Open savanna
- Woodland

**GPS**

- Cultivated areas
- Forest
- Woodland

Projection: UTM Zone 36 S  
Datum: WGS84

0 500 1,000 2,000 Km

Figure 4. Vegetation types on Mt Chiperone, supervised classification.

In the digital classification map the separation between forest and open forest is not clear owing to lack of ground control points in this area and from casual ground observation. A better figure for forest extent is probably obtained by combining these two classes. As can be seen from Table 1, this is around 1635 ha. As mentioned previously, the separation of medium and high altitude forest was based solely on the 1600 m contour.

Ground control points were limited to the southern slopes, with a few on the eastern slopes (see Figure 1). The limited distribution of these points across the mountain was a severe impediment to a more accurate determination of vegetation patterns.

#### c) *Historical Forest Cover*

The 1969 extent of moist forest cover was determined from stereoscopic visual analysis of historical air photos. The mapped extent was calculated manually using a dot planimeter as no reliable control points were available to enter it into a GIS. Allowance was made for edge-distortion by calculating areas only from the central parts of the air photos. Total area was calculated to be around 2500 ha.

#### **Historical Change**

The limited comparisons possible on the extent of forest cover as seen on historical air photos, on recent satellite imagery and on the ground suggest that forest cover has diminished by 800–1000 ha over the intervening 37 years, or between 32 and 40%. This would appear to be primarily from the southern and south-eastern slopes between 800–1200 m, in some of the northern valleys at similar altitudes, and on lower south-western slopes at 600–1000 m.

Ground observation shows that fire is used extensively for clearing fallow land before replanting, and these fires eat into the remaining forest, particularly in gullies where fires are fiercer owing to a denser combustible layer of plant matter (= fuel). Evidence for this can be seen in severely fire-damaged forest trees now standing clear of the remaining forest, and in the frequency of secondary woody species growing into the forest. Even 50 m inside the present forest margin, evidence of fire on the trunks of forest trees can be seen. Apart from on the southern and south-eastern slopes, it is possible that much of the agricultural clearance in recent years is actually of pre-existing cleared land, or land that 50 years ago or more supported woodland, rather than true moist forest.

#### **Vegetation Types**

Broadly speaking, Mt Chipirone is covered with medium and higher altitude forest above about 1000 m, with miombo or similar woodland types below that and on the ridges above with shallow soils. The topography on the mid-slopes is very dissected and often steep, hence soils are shallow.

This may be an explanation for the comparatively low number of large diameter trees. Species composition and structure of the forest gradually changes at around 1600–1800 m, with shorter, more sclerophyllous tree species festooned with 'bearded' lichen (probably *Usnea* spp.), indicating cooler conditions and more frequent moist air (i.e. mists and cloud). At the very peak an odd sclerophyllous thicket vegetation is found, insufficiently investigated or collected, comprising *Erica* (previously *Phillipia*) shrubs and the shrubby *Aloe arborescens*. The general vegetation patterns and composition are similar to those recorded from Mt Mulanje by Chapman & White (1970).

From a conservation viewpoint, the most important habitat is moist forest. Such forests are particularly extensive and relatively undisturbed on Mt Chipirone compared to many areas in adjacent Malawi. Moist forest can be subdivided into medium altitude and high altitude, with the division occurring around 1600 m. In practice there is probably a broad transition zone ranging from 1600–1800 m depending on slope and aspect. Some tree species more typical of higher

altitude forest are found in medium altitude forest, and in more sheltered and favourable positions medium altitude forest species, e.g. *Khaya anthotheca* and *Strombosia scheffleri*, can be found as high as 1800 m. From limited observations it appears the forest canopy is around 20–30 m high at 1500 m altitude, reduced to 10–20 m high at 1700 m.

Unlike some other mountains in the region (e.g. Namuli, Gorongosa, Mulanje), Mt Chipirone has no grassland, a habitat that often supports endemic species. The main open areas are patches of bare or barely vegetated or scrubby rock outcrops.

Much of the woodland below 800 m altitude has been cleared for subsistence cultivation, at least where soils are more fertile. Large patches of miombo woodland are still present on the western and northern slopes (these areas were not visited), where human settlements are very few. Much of this cleared area supports regenerating or secondary woodland and fallows. The main vegetation types are described below.

### Summit Thicket

This type is only reported from the exposed peak at around 1900–2000 m altitude to the summit, and probably has an extent of only 2–5 ha. It comprises an impenetrable thicket of *Erica* cf. *johnstoniana* shrubs 2–3 m high, previously classified under *Phillipia*, with a mass of decumbent stem aloes (*Aloe arborescens*). A small succulent, *Crassula swaziensis*, was found on rocks, along with the fern *Mohria lepigera*. The three persons from the expedition who reached this point found it too difficult to hack away through to the actual (unmarked) peak. Epiphytic "bearded" lichens (cf. *Usnea* spp.) are common.

Unfortunately, owing to time constraints, very little collecting was done here. It would appear this type is rarely, if ever, exposed to fire, and is rarely visited by humans. It is under no threat, and is of conservation interest.

### High-altitude Forest

Above about 1600 m forest species composition starts to change. Shorter and more sclerophyllous trees predominate, many festooned with lichens. Total extent is probably around 400 ha. This forest type was only superficially explored on the southern ridges, hence the description below must be considered provisional.

The main woody species appear to be *Peddiea africana*, *Diospyros whyteana*, *Maytenus undata*, *M. acuminata*, *Myrsine africana*, *Ochna holstii*, *Vepris* (*Oricia*) *bachmannii* and *Olea capensis* subsp. *macrocarpa*. Other common species include *Garcinia kingaensis*, *Tricalysia* sp., *Lasianthus kilimandscharicus*, *Psychotria zombamontana*, *Rawsonia lucida*, *Xymalos monospora*, *Tabernaemontana stapfiana*, *Schefflera goetzenii*, *Rinorea angustifolia* subsp. *ardisiifolia* and *Rapanea melanophloeos*. Various species of fern and *Selaginella kraussiana* are more common here than lower down. Trees more typical of medium altitude forest also found at 1600 m include *Diospyros* cf. *abyssinica*, *Khaya anthotheca*, *Strombosia scheffleri*, *Syzygium guineense* subsp. *afromontanum*, *Rawsonia lucida*, *Myrianthus holstii*, *Garcinia volkensii* and *Drypetes gerrardii*.

At these altitudes there are also patches of woodland and forest margins or open areas associated with rocky outcrops. Common species include *Carissa bispinosa*, *Schefflera goetzenii*, *Dovyalis macrocalyx*, *Erythroxylum emarginatum*, *Tricalysia acocantheroides* and the lithophytic fern *Oleandra distenta*.

No evidence of fire was seen, and little evidence of human use or visitation. Accessibility is difficult, and the terrain is rugged. This type is of particular conservation interest, although the extent is limited and the species are commonly found on other montane areas in southern Africa.

### **Medium-altitude Forest**

This vegetation type probably comprises the majority of the area on the mountain above 800 m altitude, although there is very little forest on the side visited below about 1000 m. Unfortunately, only the forests of the southern slopes from 1000–1200 m were adequately explored, hence medium altitude forests on other parts of Mt Chipirone may differ from the description below. The approximate extent is 1300 ha, most of it on steep slopes with very few gaps or breaks. Tree heights can reach 40–50 m in gullies and other favourable localities, but generally trees are 20–30 m high. Canopy height decreases with increasing altitude and on more shallow soils. The stocking rate of large diameter trees is lower than in some other regional medium altitude forests.

The main tree species are *Newtonia buchananii* (particularly prominent on ridges), *Strombosia scheffleri* (with purplish flaking bark), the thinner-stemmed *Rinorea convallarioides*, fluted trunks of *Chrysophyllum gorungosanum* and scattered large individuals of *Khaya anthotheca*, some of them quite magnificent. There are very few emergents. Large strangling *Ficus* trees are few and scattered throughout. Other common trees include *Rothmannia urcelliformis*, *Drypetes gerrardii* and *Myrianthus holstii*, while somewhat smaller or sub-canopy trees include the large-leaved *Funtumia africana*, *Rawsonia inermis*, *Rinorea ferruginea*, *Garcinia kingaensis*, *G. volkensii*, *Trilepisium madagascariense* and *Pleiocarpa pycnantha*. The understorey vegetation is not thick and is characterised by *Dracaena fragrans* and *Pseuderanthemum subviscosum*, along with scattered forest grasses and the scrambling *Behnia reticulata*. There are a lot of young regenerating plants and seedlings of *Chrysophyllum* and *Rinorea convallarioides*. Lianas such as *Agelaea pentagyna* are not very common.

Where the forest adjoins woodland or land cleared for agriculture, typical forest edge species are *Albizia* cf. *gummifera*, *Macaranga capensis* and *Trema orientalis*, with large individuals of *Englerophytum magalismontanum* and *Parinari excelsa* on the ridges.

The three main features of conservation interest are (i) the extensive area of forest found, (ii) the uninterrupted altitudinal sequence it covers from 1000 to 2000 m, and (iii) the relatively intact and undisturbed nature of the forest. There has been some minor tree-felling close to the forest margins, but the biggest threat is uncontrolled fire from field clearance eating into the forest, especially in the gullies. There is evidence of such fires severely damaging or destroying even large forest trees up to 50 m from the margins, with an associated loss of humus from the forest floor and destruction of the low shrub and herbaceous layers. Once this happens, a number of forest gap or edge species such as *Trema* and *Albizia* establish themselves inside the forest along with a thick herbaceous undergrowth, which can inhibit regeneration of forest trees through excessive competition.

### **Miombo and Similar Woodland Types**

Miombo woodland is seasonally-deciduous with canopy cover ranging from 20–80% comprising trees of *Brachystegia* and/or *Julbernardia*, and with a well-developed grass layer underneath. On Mt Chipirone such woodland is mainly characterised by various *Brachystegia* species, but there are also some areas without *Brachystegia* but with other miombo-associated species.

The woodlands studied were at an altitude of 600–1100 m on the southern and south-eastern flanks. At lower altitudes woodlands have often been cut and/or frequently burnt, but were in a better condition at altitudes above 900 m. At 600–800 m woodland covered the slopes of ridges as well as the ridge-tops, with the less steep land having been cleared for cultivation, while at higher altitudes

woodland was mostly confined to the shallow soil ridges, sometimes almost surrounded by moist forest. Various woodland types were noted. Most ridges were dominated by *Brachystegia spiciformis*, whilst others supported *B. utilis* or *B. tamarindoides* subsp. *microphylla* (= *B. glaucescens*), while one ridge at 1200 m had an open woodland of *Acacia abyssinica*. *Brachystegia boehmii* and some typical miombo species (e.g. *Schrebera trichoclada*, *Securidaca longepedunculata*, *Ozoroa reticulata*, *Elephantorrhiza goetzei*) were only found at an altitude of 650 m or below.

The small extent of 5–10 m high *Acacia abyssinica* woodland was unusual and seemed to be confined to just one ridge. Associated species included the fern *Pteridium aquilinum*, a scrambling *Rubus* sp. and with *Trema orientalis* at the forest/woodland boundary. Of particular note was the high number of epiphytic ferns, orchids and lichens on the *Acacia* branches, while the root parasite *Sarcophyte sanguinea* subsp. *piriei* was very common and visible on the ground underneath. It was flowering extensively in late November, appearing above ground in blackish-red masses.

*Brachystegia spiciformis*-dominated woodland with a 10–15 m high canopy also contained miombo species such as *Bridelia micrantha*, *Combretum molle*, *Pericopsis angolensis*, *Pterocarpus angolensis*, *Erythrina abyssinica*, *Cussonia arborea*, *Uapaca kirkiana*, *Uapaca nitida*, *Psorospermum febrifugum*, *Parinari excelsa*, *Faurea saligna*, and some typical forest edge species such as *Harungana madagascariensis* and *Albizia gummifera*. Epiphytic orchids and ferns were also very common on the branches. It does not appear as if these areas previously supported forest, at least over the last 50–100 years.

Although most wooded ridges were dominated by *B. spiciformis*, others were somewhat drier or dominated by trees of *Brachystegia utilis* 8–10 m high, owing to more shallow, less moisture-retentive soils. Associated species were *Julbernardia globiflora*, *Uapaca kirkiana*, *Protea welwitschii*, *Pterocarpus angolensis* and *Monotes africana*, with one patch of the fine-leaved *Brachystegia tamarindoides* subsp. *microphylla*.

The upper reaches of gullies below the ridges mostly support regenerating woodland and/or forest with secondary species such as *Trema orientalis*, *Macaranga capensis* and *Bridelia micrantha*. These gullies may well have supported some type of forest previously, now destroyed by fire.

Although interesting, such woodlands and the species comprising them are very widespread across this part of Africa. They have a useful role in forming part of the forest–woodland–shrubland vegetation mosaic on the mountain and acting as a buffer to the forest itself, and there are undoubtedly a number of plant, vertebrate and invertebrate species that on Chipirone are only found here.

### **Disturbed Woodland and Fallows**

Most of this vegetation type was found below 900 m altitude and in the lower reaches of gullies between ridges. In most cases it is probable that it was previously (5–50 years ago) forest or forest-edge vegetation, but has been destroyed by frequent fires and perhaps partial clearance. This type was not investigated or collected in any detail owing to its low conservation value and potential.

In the gullies vegetation could be quite thick, making progress difficult. In more recent fallow it was more open, and a number of areas had been cleared within the last two years. Characteristic trees included *Trema orientalis*, *Dombeya burgessiae*, *Bridelia cathartica*, along with the plants of lower height such as *Smilax anceps*, *Annona senegalensis*, *Fromomum* spp., *Mucuna pruriens*, clumps of bananas (*Musa*) and the bamboo *Oxytenanthera abyssinica*. The liana *Entada rheedei* with characteristic large woody pods was found in some places.

Lower down the slopes, at around 800 m and below, significant areas of bamboo (*Oxytenanthera abyssinica*) thicket were found. These appear to develop after clearance for cropping, particularly in gullies, and are tolerant of repeated fire.

Also at lower altitudes (c.600 m) a riparian forest fringe can be found along larger watercourses. Although extensively cleared in places, remnant large trees still remain, including *Treulia africana*, *Synsepalum* (= *Pachystela*) *brevipes*, and *Ficus* species.

### Rocky Outcrops

There were a few areas of bare, or almost bare, rock on the mountain, but only two or three were visited owing to access problems. From airphotos there appears to be a greater extent of such outcrops on the north and north-eastern part of the mountain.

Surprisingly, even the rugged and narrow ridges have woody vegetation on them. Where rock is exposed over an area, vegetation consists of low shrubs, *Aloe*, and grasses/sedges, typical of such environments elsewhere. Through binoculars it appeared that some rock outcrops had been burnt. Further information is not available.

## 5. BOTANY

Plants were collected fairly extensively in the area around the main Forest Camp on the southern flanks of Mt Chipirone (Figure 1 shows collecting points) and along the track to the village below. More limited collecting was done above Marega village on the eastern side. In addition, plant specimens were collected from the summit ridge at altitudes of 1500 and above, which proved to be most interesting as vegetation composition appears to change above 1600–1800 m.

Most specimens were collected with notes on locality and habit. Unfortunately, time did not allow for good notes to be recorded from collections on the summit ridge. Labelled specimens are deposited at the LMA Herbarium at IIAM in Maputo and at the Kew Herbarium in London. A partial third set is deposited in Zomba (MAL). A total of over 400 numbered and unnumbered specimens were recorded.

A species list compiled from both specimens and confirmed field sightings is given as Annex 2. The great majority of records are from above 800 m altitude, although some woodland species from below this altitude are included. Of the 229 taxa listed, 145 are woody plants (trees, shrubs, lianas).

A number of species (around 15) are not listed in the Sabonet Mozambique plant checklist (Da Silva, Izidine & Amude 2004). However, this checklist is incomplete and only represents collections in the National Herbarium in Maputo. When citations from families published in the Flora Zambesiaca are incorporated, virtually all taxa are recorded from Mozambique, the great majority being recorded from the Z, MS or N divisions. However, there may be some new taxa for Mozambique in as yet unpublished family treatments. It should also be noted that northern Mozambique is known to be poorly and patchily collected, part of the justification for the present study.

Species or points of particular notes include:

*Widdringtonia whytei* — although common on Mt Mulanje, this tree was not seen on Chipirone.

*Aloe arborescens* (Aloaceae) – thicket-forming stem *Aloe* on rock outcrops at higher altitudes and on the peak. Very local here, but common on Mt Mulanje.

*Cyperus amauropus* (Cyperaceae) — first record for Mozambique.

*Dracaena fragrans* (Dracaenaceae) — first record for Mozambique Z.

*Pollia condensata* (Commelinaceae) – common forest undergrowth herb. First record of the genus from the FZ area.

*Abrus melanospermus* (Fabaceae: Papilionoideae) — first record for Mozambique.

*Coffea mufindensis* (Rubiaceae) — regarded as Vulnerable in the Mozambique Red Data List.

*Crassula swaziensis* (Crassulaceae) — this species has a number of varieties, one of which is apparently confined to the Namuli massif in N Mozambique. Material was insufficient for further determination, but this collection may well extend the known range of what was considered a Namuli endemic.

*Plectranthus kapatensis* (Lamiaceae) — first record for Mozambique Z.

*Strychnos* sp. (Loganiaceae) – a forest liana not possible to match at Kew; a potentially interesting taxon.

Three species of *Rinorea* were found in the forest (*R. convallarioides*, *R. ferruginea*, *R. angustifolia* subsp. *ardisiiflora*) compared to only one (Strugnell 2006) or two (White, Dowsett-Lemaire & Chapman 2001) recorded from the more extensive forests on Mt Mulanje. It is not known why this should be. *R. convallarioides* is common at around 1000–1200 m, while *R. angustifolia* appears to be more common at altitudes above 1600 m. *Rinorea* is primarily a genus of smaller trees from coastal or lowland forests, which may suggest Chipirone has more lowland or coastal influence than Mt Mulanje. This lowland and 'East African coastal' influence can also be seen in the presence of *Funtumia africana* and *Englerophyllum natalense* and *Aporrhiza paniculata*, and, on the lower slopes, *Treulia africana* and *Synsepalum brevipes*. In Zimbabwe, all *Rinorea* species are considered Critically Endangered as they are only found in small remnant patches of low-altitude forest.

The only recorded species from Chipirone in a formal threat category (Vulnerable, Endangered or Critically Endangered) for Mozambique in the Sabonet Red Data List (Izidine & Bandiera 2002) is *Coffea mufindensis*. However, some species found are considered threatened in Zimbabwe as they are low altitude forest species there confined to small remnant forest patches along the Haroni and Rusitu rivers, but are more widespread in adjacent parts of Mozambique.

## 6. ZOOLOGY

Despite the close proximity and accessibility of Mt Chipirone to Mt Mulanje in Malawi considerably less biological work has been undertaken into the biodiversity of this area compared to the large volume of work undertaken on Mt Mulanje. There have been two previous visits to assess aspects of the biology of the area. The first was by Mr Jali Makawa, a bird collector for C.W. Benson, who made a 6-day collecting expedition in 1950 to Mt Chipirone as part of a wider birding survey of montane areas in Zambézia province (Benson 1950). Since then the only documented



visit was made in November 2005 by Claire Spottiswoode, Hassam Patel, Eric Herrmann and Julian Bayliss in preparation for this current Darwin expedition. During this latter trip the threatened avifauna was assessed and some plants collected. There are unconfirmed reports of an earlier visit (around 2002) by the American Field Museum of Natural History to collect birds and small mammals, but no results are available.

Findings from the present expedition show that generally the moist forest area was relatively undisturbed with no evidence of logging. However, there was much evidence of man-made fires on ridges, possibly as a result of preparation for small scale cultivation. The forest area above 1000 m is hunted with the main method being gin trapping. The main animals hunted are Bushbuck *Tragelaphus scriptus*, Bushpig *Potamochoerus larvatus*, and a duiker *Cephalophus* spp. Gin traps are apparently commonly available in local markets. The Leopard *Panthera pardus* is reputed to be common throughout the forest. Of particular note was the mention on several occasions of the presence of Buffalo *Syncerus caffer* on the northwestern side of Mt Chipirone. The forest is also utilised for honey collecting.

A relatively high incidence of calling bushbabies was recorded from around the main forest camp at c.1000 m. The calls sounded similar to those of the Lesser Bushbaby that occurs on Mt Mulanje, which has been identified as *Galago grantii*. This would merit further investigation.

This study found two endemic species previously only known to occur on Mt Mulanje in Malawi, 60 km to the north, one lizard and one butterfly. This is not wholly surprising as the distance is relatively small, but such records represent important new discoveries for Mozambique.

### Local Hunter Records

One of the forest guides employed on the expedition also utilised the area for hunting. Employing local hunters as forest guides is a good practice as they are often the most knowledgeable people on the area and know where various ground traps have been set, traps that can severely injure humans.

Table 2. Mammals recorded as being found on Mt Chipirone.

Common Name	Scientific name	Local name (in Khokhola)	Notes
Bushpig	<i>Potamochoerus larvatus</i>	Nguluwe	
Blue Monkey	<i>Cercopithecus mitis</i>	Nchimwe	
Vervet Monkey	<i>Cercopithecus aethiops</i>	Nakahmwa	Black face. Woodland
Baboon	<i>Papio cynocephalus</i>	Nyani	Woodland
Duiker	<i>Cephalophus</i> spp	Nazoro	Black body, reddish head
Rock Hyrax	<i>Procavia capensis</i>	Mbila	
Bush baby	<i>Galago granti?</i>	Nchanga	
Bushbuck	<i>Tragelaphus scriptus</i>	Nanse	
Porcupine	<i>Hystrix africaeaustralis</i>	Nansununga	Forest and woodland
"Big Rat"	?	Ncheza	White and black spots, small tail twice length of back leg
Black Rat	<i>Rattus rattus?</i>	Nyenga	Same size as Ncheza but with 30 cm tail
Mole Rat		Nafoko	Brown/grey back, white belly. No tail. Woodland. Lives underground
Zorilla	<i>Ictonyx striatus</i>		Rat-like teeth, round small ears

The most common form of trapping in Mt Chipirone is 'gin-trapping' (also know as Bear Traps in the western world), not snares or the setting of fires to flush animals into traps as is the case on Mt Mulanje. A local hunter and guide, Besta, was interviewed on the animals present in the forest and which were commonly hunted by the local population. His responses are given in Table 2.

### Small Mammals

Small mammal species were also opportunistically collected. Such studies largely centred around the mist netting of bats. Although time was very limited, several species were caught (Table 3). The capture of *Miniopterus inflatus* represents an important new record for the region. As can be seen from the distribution map (Figure 5) it has a very restricted range and is only known from a few localities. It was previously unknown from Mozambique and was only recently caught at Mt Gorongosa (Ara Monadjem, pers. comm.), about 300 km south of Chipirone. The Chipirone record is only the second for the country.

Table 3. Small mammals collected from Mt Chipirone (identified by Peter Taylor, Durban Natural History Museum).

Species	Sex	Comments
<i>Miniopterus inflatus</i>	F	*Greatest skull length (CIL=16.0)
<i>Myotis tricolor</i>	F	*CIL. 6 molars top and bottom
<i>Myotis tricolor</i>	F	*CIL. 6 molars top and bottom
<i>Praomys delectorum</i>		




Figure 5. Known records for *Miniopterus inflatus* in Southern Africa (Peter Taylor, 2007).

### Birds

The avifauna of northern Mozambique is very poorly known, and many montane areas are still largely unexplored for birds except from single collecting expeditions between 1932 and 1950 (e.g. Vincent 1933a, 1933b, 1934). Consequently there have been repeated calls for further descriptive information on the extent, conservation status and avifauna of the evergreen forests of this region (Collar & Stuart 1988, Stattersfield et al. 1998, Parker 2001), with particular focus on the status of the rapidly declining Thyolo Alethe *Alethe choloensis* (BirdLife International 2006).

Mt Chipirone was visited 15–18 December 2005 by Claire Spottiswoode, Hassam Patel, Eric Herrmann and Julian Bayliss in preparation for this expedition in November 2006 (Spottiswoode et al. 2007, in press). Bird species were detected by sightings, with cassette and minidisc recordings sometimes subsequently used for playback, and by mist-netting. Mist-nets were opened for a total of 56 net-hours. To investigate the forest areas a camp was made at 1050 m; the highest altitude reached was 1260 m. Incidental observations were also made while hiking through mixed

woodlands and cultivation at lower altitudes (500–1000 m). The resulting checklist is given in Annex 3.

According to Spottiswoode (pers. comm.) the two globally threatened species collected by Makawa in 1950 were both found in 2005. A territorial pair of Thyolo Alethe *Alethe choloensis* (Endangered) were seen and tape-recorded in ridge forest at altitude 1200 m (16°30'12"S, 35°43'50"E), and a pair of White-winged Apalis *Apalis chariessa* (Vulnerable) were seen in a forest clearing at altitude 1120 m (16°30'20"S, 35°43'50"E).

Further to these findings a considerable range extension was represented by the numerous individuals of Eastern Bronze-naped Pigeon *Columba delegorguei*, heard constantly at 1100–1200 m and seen pursuing territorial chases. This is the only record between Zimbabwe's Eastern Highlands and central Tanzania, other than the handful of records from Thyolo Mountain in Malawi (Dowsett-Lemaire & Dowsett 2006), where it is now likely to be extinct in view of virtual complete deforestation of this area. The fact that Makawa did not find this species in 1950 is not surprising. He visited the area in the non-breeding season (July) when this species, if not calling, would have been very inconspicuous (Benson & Irwin 1966).

In addition to these montane species, other species typical of forest margins, miombo and mixed woodland and *Pennisetum* grassland habitats were recorded, such as the Olive-headed Weaver *Ploceus oliviceps* and Cabanis' Bunting *Emberiza cabanisi* (miombo), and Marsh Tchagra *Tchagra minuta* and Singing Cisticola *Cisticola cantans* (grassland).

Occurrence of globally threatened, biome-restricted, and/or range-restricted species were noted. Table 4 shows globally threatened species in bold type, along with their 2005 threat status. Biome and Endemic Bird Area (EBA) membership for biome-restricted and range-restricted species (Fishpool & Evans 2001) is indicated by the following: A07 Afrotropical Highlands Biome; A09 East African Coast Biome; A10 Zambesian Biome; 105: Tanzania-Malawi Mountains EBA. Records for Mt Chipero from 1950 are from Benson (1950).

Table 4. Globally threatened, biome-restricted, and/or range-restricted bird species from Mt Chipero (from Spottiswoode et al., in press).

Species	Biome	EBA	Chipero 1950	Chipero 2005
Bar-tailed Trogon <i>Apoloderma vittatum</i>	A07		x	
Grey Cuckooshrike <i>Coracina caesia</i>	A07		x	
Striped-cheeked Greenbul <i>Andropadus milanjensis</i>	A07		x	
Orange Ground-thrush <i>Zoothera gurneyi</i>	A07		x	
<b>Thyolo Alethe <i>Alethe choloensis</i></b> EN	A07	105	x	x
White-starred Robin <i>Pogonocichla stellata</i>	A07		x	
<b>East Coast Akalat <i>Sheppardia gunningi</i></b> VU	A09			
Olive-flanked Robin-chat <i>Cossypha anomala</i>	A07		x	
<b>White-winged Apalis <i>Apalis chariessa</i></b> VU	A09	105	x	x
Yellow-throated Warbler <i>Phylloscopus ruficapilla</i>	A07		x	x
White-tailed Crested Flycatcher <i>Elminia albonotatus</i>	A07		x	
Green-headed Oriole <i>Oriolus chlorocephalus</i>	A09		x	x
Eastern Double-collared Sunbird <i>Nectarinia mediocris</i>	A07		x	
East African Citril <i>Serinus hypostictus</i>	A07			x
Red-faced Crimson-wing <i>Cryptospiza reichenovii</i>	A07		x	
Swee Waxbill <i>Estrilda melanotis</i>	A07			x
Bertram's Weaver <i>Ploceus bertrandi</i>	A07			x
Olive-headed Weaver <i>Ploceus oliviceps</i>	A10			x

## Reptiles and Amphibians

Reptiles and amphibians were collected opportunistically but non-intensively during the expedition. Only a few records have been collated for Mt Chipirone, but which contain some notable and interesting results (Table 5).

Of particular note was the observation record of the Gaboon Viper (*Bitis gabonica*), seen on the forest margin on 25 November 2006. It is the first record of this species from Mt Chipirone and the first record from this region of Mozambique since 1950. The earliest record from northern Mozambique, and apparently the only voucher specimen from this half of the country (Don Broadley, pers. comm. 2007), is from Wilhelm Peters in 1846 (published in *Reise nach Mossambique*: 146, as *Bitis rhinoceros*) from Prazo Boror (c.100–200 m altitude), just south of Morrumbala Mountain and about 140 km SE of Mt Chipirone. The Gaboon Viper has not been recorded from nearby Mt Mulanje, despite the presence of apparently suitable habitat. In Malawi it has only been ever recorded from the Mzuzu–Nkhata Bay area at 500–600 m altitude, almost 600 km away (Don Broadley, pers. comm.).

Table 5. Reptiles recorded from Mt Chipirone.

<b>Lizards</b> (Order Lacertilia)	
Gekkonidae	
King Dwarf Day Gecko	<i>Lygodactylus rex</i>
Chamaeleonidae	
Pygmy Chameleon	<i>Rhampholeon champmanorum</i>
<b>Snakes</b> (Order Serpentes)	
Typhlopidae	
Blunt Blind Snake	<i>Letheobia obtusus</i>
Colubridae	
Mulanje Water Snake	<i>Lycodonormorphus mlanjensis</i>
Mozambique Twig Snake	<i>Thelotornis mossambicanus</i>
Viperidae	
Gaboon Viper	<i>Bitis gabonica</i>
<b>Frogs</b> (Order Anura)	
Arthroleptidae	
Dwarf Squeaker	<i>Arthroleptis xenodactyloides</i>

Another notable record was the capture of the gecko *Lygodactylus rex*. This species has only recently been discovered on Mt Mulanje in Malawi, to which it was previously thought to be endemic. Its discovery on Mt Chipirone is the first record for Mozambique (Bill Branch, pers. comm.).

## Lepidoptera

The Lepidoptera of Mt Chipirone were opportunistically collected over the course of the expedition by Julian Bayliss. The majority of specimens were collected with a 4-fold hand net or (for the Charaxinae and certain Satyridae) through baited aerial traps using fermenting bananas.

A total of 56 butterfly species were collected and were sent to the African Butterfly Research Institute (ABRI) in Nairobi, Kenya for formal identification. ABRI is the recognized institute where the main African butterfly reference collection is stored. There appear to be no previous records of Lepidoptera from Mt Chipirone, and therefore the list in Table 6 presents the first records from this site.

The range of butterflies typically represents a wet forest, forest edge and miombo woodland collection. Of particular note was the capture of *Cymothoe melanjae* (an endemic previously only known from Mt Mulanje), *Eurema senegalensis*, *Eurema floricola*, *Bicyclus vansoni*, *Anthene lunube* and *Platylesches vasta*. These are the first records of these species from Mozambique (Steve Collins, pers. comm.).

Table 6. List of butterflies collected from Mt Chipirone (Nov/Dec 2006), identified by J. Bayliss and S. Collins (ABRI).

Family/subfamily	Species/authority
<b>Hesperiidae</b>	
Hesperiinae	<i>Acada bieriatius</i> Mabille 1893
Hesperiinae	<i>Platylesches rasta</i> Holland 1896
Pyrginae	<i>Eagris sabadius</i> Lathy 1901
Pyrginae	<i>Tagiades flesus</i> Fabricius 1781
<b>Lycaenidae</b>	
Lipteninae	<i>Alaena amazoula</i> Hawker Smith 1933
Lipteninae	<i>Teriomima puella</i> Kirby 1887
Polyommatae	<i>Anthene barnesi</i> Stevenson 1940
Polyommatae	<i>Anthene lunulata</i> Doubleday 1847?
Polyommatae	<i>Euchrysops malathana</i> Boisduval 1833
Polyommatae	<i>Leptotes pirthous</i> Linnaeus 1767
Theclinae	<i>Axiocerces</i> sp.
<b>Nymphalidae</b>	
Acraeinae	<i>Acraea acrita</i> Hewitson 1865
Acraeinae	<i>Acraea calderena</i> Hewitson 1877
Acraeinae	<i>Acraea egina areca</i> Cramer 1775
Acraeinae	<i>Acraea johnstoni</i> Godman 1855
Acraeinae	<i>Acraea natalica</i> Boisduval 1847
Acraeinae	<i>Acraea oncaea</i> Hopffer 1855
Acraeinae	<i>Acraea zetes</i> Linnaeus 1758
Argynninae	<i>Lachnoptera ayresii</i> Trimen 1879
Biblidinae	<i>Eurytela hiarbas</i> Rothschild & Jordan 1903
Charaxinae	<i>Charaxes brutus natalensis</i> Staudinger & Schatz 1886
Charaxinae	<i>Charaxes candiope</i> Godart 1824
Charaxinae	<i>Charaxes cithaeron nyassae</i> van Someren 1964
Charaxinae	<i>Charaxes protoclea azota</i> Hewitson 1877
Charaxinae	<i>Charaxes pollus geminus</i> Rothschild & Jordan 1900
Charaxinae	<i>Charaxes violetta melloni</i> Fox 1963
Charaxinae	<i>Euxanthe wakefieldi</i> Ward 1873
Danainae	<i>Amauris niavius dominicanus</i> Trimen 1879
Limnitiinae	<i>Cymothoe melanjae</i> Bethune-Baker 1926
Limnitiinae	<i>Neptis alta</i> Overlaet 1955
Limnitiinae	<i>Neptis saclava</i> Hopffer 1855
Limnitiinae	<i>Neptis swynnertoni neavi</i> Trimen 1912
Nymphalinae	<i>Junonia artaxia</i> Hewitson 1864
Nymphalinae	<i>Junonia natalensis</i> Felder & Felder 1860
Nymphalinae	<i>Antanartia schaenia dutia</i> Howarth 1966
Nymphalinae	<i>Precis antilope</i> Feisthamel 1850
Nymphalinae	<i>Precis tugela</i> Trimen 1879
Nymphalinae	<i>Salamis parhassus</i> Drury 1782
<b>Papilionidae</b>	
Papilioninae	<i>Graphium angolanus</i> Goeze 1779

Papilioninae	<i>Graphium polices</i> Cramer 1775
Papilioninae	<i>Papilio dardanus tibullus</i> Kirby 1880
Papilioninae	<i>Papilio echeriodes shirensis</i> Hancock 1987
Papilioninae	<i>Papilio nireus</i> Doubleday 1845
<b>Pieridae</b>	
Coliadinae	<i>Catopsilia florella</i> Fabricius 1775
Coliadinae	<i>Eurema floricola</i> Boisduval 1883
Coliadinae	<i>Eurema regularis</i> Butler 1876
Coliadinae	<i>Eurema senegalensis</i> Boisduval 1836
Pierinae	<i>Appias sabina phoebe</i> Butler 1901
Pierinae	<i>Mylothris sagalla</i> Butler 1896
Pierinae	<i>Nepheronia thalassine</i> de Boisduval 1836
<b>Satyridae</b>	
Satyriinae	<i>Bicyclus safitza</i> Westwood 1850
Satyriinae	<i>Bicyclus vansoni</i> Condamin 1965
Satyriinae	<i>Gnophodes betsimena diversa</i> Butler 1880
Satyriinae	<i>Henotesia (Heteropsis) perspicua</i> Trimen 1873
Satyriinae	<i>Melanitis leda</i> Westwood 1851
Satyriinae	<i>Melanitis libia</i> Distant 1882

Several day-flying moths were also captured and most await identification. The capture of the geometrid moth *Cartaletis nigricosta* is an indicator of much wetter forest. It was originally described from Mt Mulanje which was the only known locality until specimens were collected from Mt Rungwe (Tanzania) and now Mt Chipirone (Herman Staude, pers. comm.). Larvae of a moth attacking *Khaya* seedlings, probably *Hypsipyla robusta*, were later found hatching from seeds collected on the lower slopes of Mt Chipirone (T. Alves, pers. comm.).

### Coleoptera and Hemiptera

The following are some preliminary identifications by Cornell Dudley of Coleoptera (beetles), Hemiptera (bugs) and Diptera (flies) collected during the expedition to Mt Chipirone.

#### Coleoptera: Cetoniidae

<i>Amaurodes passerinii</i> Westwood	common
<i>Rhabdotis aulica</i> Fabricius	very common
<i>Diplognatha silicea</i> MacLeay	very common
<i>Plaesiorrhina mohondana</i> Oberthür	uncommon
<i>Stethopseudinca maculatus</i> Valak Lucassen	rare, previously only known from Mt Mulanje

#### Coleoptera: Scarabaeidae-Coprinae

<i>Diostellopalpus neavei</i> d'Orbigny	common, but previously only known from Mulanje
---	--

#### Coleoptera: Rutelidae

<i>Anomala</i> sp.	large difficult genus
--------------------	-----------------------

#### Coleoptera: Cerambycidae-Lamiinae

<i>Tragocephala pretiosa</i> Hintz	rare
------------------------------------	------

#### Coleoptera: Chrysomelidae-Cassidinae

<i>Aspidomorpha</i> sp.	common, but unable to identify
-------------------------	--------------------------------

#### Hemiptera-Homoptera: Cicadidae

<i>Ioba leopardina</i> (Distant)	very common
? <i>Taipinga</i> sp.	uncommon in collections, genus unsure

#### Diptera: Tabanidae

<i>Tabanus</i> sp.	Common large genus; species unlikely to be new.
--------------------	---

## 7. THREATS AND CONSERVATION ISSUES

Mt Chipirone is under no form of formal conservation and has no specific status. However, the forest and its wildlife should be protected under the Forest and Wildlife Act (Lei No. 10/99 of 1999). The Environmental Act (Lei No. 20/97 of 1997) prohibits cultivation of annual crops on slopes greater than 7° and perennial crops on slopes greater than 14°, which many of Chipirone's slopes exceed.

Both Mount Namuli and Chipirone are recognised as Important Bird Areas (MZ 009 and MZ 010 respectively) by Parker (2001). A brief description of the area is given there along with a list of the forest bird species of interest – *Alethe choloensis*, *Apalis chariessa* (only known site in Mozambique), and the woodland species *Nectarinia shelleyi*. The conservation status of these bird species is fundamentally determined by the extent and condition of the forest habitat.

At the base of the southern and eastern flanks, there are a few villages with a population around 1000–2000 people. However, there is no clean water supply other than the small streams coming off the mountain. The nearest wells or boreholes are some kilometres away. From an interview with Chief Sapande Marega at Marega locality on the eastern slopes of Mt Chipirone it appears that the forests on the upper slopes have little perceived value to the local population. The only persons going there are bushmeat hunters, a small select group. Even traditional medicinal practitioners tend to collect medicinal plants from the disturbed and woodland areas lower down.

During the course of this interview, the Chief mentioned that there was said to be "tea" plants growing on the lower northern slopes of the mountain, but it is likely this does not refer to the tea of commerce. There has been minimal extraction of timber from the forests, only from the woodlands around where *Pterocarpus angolensis*, *Lonchocarpus capassa* and *Pericopsis angolensis* have been harvested on a small-scale.

Interestingly, he also pointed out that people clear shambas (small fields) and grow maize and beans on the steep slopes with shallow rocky soils because in bad rainfall years these fields ensure at least a small harvest. In good rainfall years, fields below the mountain have a much higher production, but in times of drought fields on the mountain slopes provide some food. This is probably related to incoming moist air and reduced evapotranspiration on the middle slopes compared to the hot dry plains.

In Malawi, folk history links Mt Chipirone with Mt Mulanje through the advent of the 'chipirone' weather front that appears to start on Chipirone and move across into Malawi. This misty weather is important for the tea plantations as it brings moisture during the dry season. A good case can be made for linking conservation of the two mountains in a form of transfrontier initiative. In both cases the key issue is moisture and water, which relies on forest cover for its generation and continued supply.

The suggestion for transfrontier cooperation has already been discussed between the Mulanje Mountain Conservation Trust in Malawi and the District Administrator of Milange District in Mozambique. The District Administrator has expressed great interest in getting technical cooperation and assistance for forest conservation and control of natural resource degradation in his area. It is intended to discuss the issue further with the Provincial Governor and provincial authorities in Quelimane during 2007.

## 8. RECOMMENDATIONS

From a conservation perspective, although Mt Chipirone has threats to its habitats and biodiversity, these are generalised, and not specific at any particular species or habitat. The threats to the forest above 1000 m appear to be primarily from: (a) encroachment by clearance for cultivation (localised, but are a particular concern on the southern and eastern slopes, and (b) from wild fires eating into the forest margin, particularly in gullies. The following specific conservation recommendations are therefore given:

- Conserve the remaining areas of moist forest, especially those above the 1000 m contour. Stop wherever possible any clearance of vegetation above this line, or perhaps even as low as 900 m.
- Control wildfires along the margins of moist forest. Fires resulting from bush clearance on steep slopes, especially gullies, burn up into the forest and destroy younger trees and the regenerating layer. Subsequently, soils do not appear to retain as much moisture, and the areas are invaded by secondary or forest margin species.
- Transfrontier cooperation and conservation initiatives with MMCT and others in Malawi should be actively encouraged. Issues of forest conservation should be linked in to water supply and rainfall. This could be initiated through a meeting with the Governor of Zambézia Province.
- A meeting should be arranged between the Provincial Director of Agriculture, the Milange District Administrator, MMCT and provincial representatives of IIAM, to determine how conservation of Mt Chipirone and the improvement of agricultural practices could best be tackled.
- Experience of catastrophic landslips on Mt Tumbine above Milange town owing to deforestation, should be used as an illustration of what unregulated forest destruction can do.

### Research Issues

- Investigate why woodlands and forests of western and northern slopes, which appear more intact although perhaps drier, are more intact than those on southern and eastern slopes.
- Determine what the altitudinal transition point of medium and higher altitude forest types is, and whether this transition is similar on all sides of the mountain.
- Further botanical and ornithological survey work is required on moist forest areas on the north-western and eastern slopes of the mountain. Further collecting is also needed in the areas of shrubland and rocky outcrops, both of which are likely to support some interesting species and outlying populations.

## 9. CONCLUSIONS

Mt Chipirone is a steep-sided montane massif that is relatively undisturbed above c. 1000 m altitude up to the summit at just over 2000 m. Owing to its conical shape, the area above 1800 m is quite small. The wooded slopes, many of which have been cleared for agriculture on the southern and eastern sides, give way to moist forest at around 1000 m altitude on the steeper slopes up to the summit. The forest is well-developed, is around 1700 ha in extent (estimates from 1635 to 1720),



and has an uninterrupted altitudinal sequence from medium altitude to high altitude. This feature is increasingly rare to find on montane massifs in the region owing to encroachment on lower slopes. The exposed summit ridge supports a fynbos-type scrub vegetation of very limited extent, and small patches of miombo-type woodlands are found on the ridges from 800–1200 m.

Owing to difficulties with access, the study was primarily confined to the southern and eastern slopes up to 1200 m. It is possible that findings may differ significantly on the northern slopes, and conservation conclusions should be interpreted accordingly.

The plant species found are mostly fairly widespread across other Eastern African mountains from South Africa through Zimbabwe and Mozambique to Malawi and Tanzania. Although a few new records for N Mozambique were recorded, no threatened species or species of particular interest or concern were noted.

Mt Chipirone supports perhaps the largest known population of the Thyolo Alethe, a threatened bird species, which is under increasing threat in Malawi owing to forest destruction. A number of interesting new records for birds, small mammals, reptiles and butterflies were also noted, making the mountain of particular conservation interest and concern for Mozambique.

## 10. REFERENCES

- Barbosa, L.A.G. (1952). Esboço da Vegetação da Zambézia. Separate from Documentário Moçambique No. 69. Centro de Investigação Científica Algodoeira, Lourenço Marques.
- Bayliss, J. (2006). Trip Report, Mount Chipirone Expedition, 22 November–5 December 2006. MMCT, Mulanje.
- Benson, C.W. (1950). A collection from Chipironi Mountain, Portuguese East Africa. *Bulletin British Ornithological Club* **70**: 51.
- Chapman, J.D. & White, F. (1970). *The Evergreen Forests of Malawi*. Commonwealth Forestry Institute, Oxford.
- Collar, N.J. & Stuart, S.N. (1985). *Threatened Birds of Africa and Related Islands*. International Council for Bird Preservation, Cambridge.
- Da Silva, M.C., Izidine, S. & Amude, A.B. (2004). A preliminary checklist of the vascular plants of Mozambique. Southern African Botanical Diversity Network Report No.30. SABONET, Pretoria.
- Dixey, F. (1927). The Mlanje Mountains of Nyasaland. *Geographical Review* **17**: 61–626.
- Dowsett-Lemaire, F. (1989). Ecological and biogeographical aspects of forest bird communities in Malawi. *Scopus* **13**: 1–80.
- Eastwood, F. (1988). *Guide to the Mulanje Massif* (3rd edition). Lorton Communications, Johannesburg.
- Izidine, S. & Bandiera, S.O. (2002). Mozambique. In: *Southern African Plant Red Data Lists* (edited by J.S. Golding), pp. 43–60. SABONET, Pretoria.

- Kassam, A.H., Van Velthuis, H.T., Higgins, G.M., Christoforides, A., Voortman, R.L. & Spiers, B. (1981). Climatic data bank and length of growing period analysis. Field Document No. 33, Project MOZ/75/011, Assessment of Land Resources for Rainfed Crop Production in Mozambique. FAO, Rome.
- Parker, V. (2001). Mozambique. In: *Important Bird Areas in Africa and Associated Islands: Priority sites for conservation* (edited by L.D.C. Fishpool & M.I. Evans), pp. 627–638. Pisces Publications/BirdLife International, Newbury & Cambridge.
- Pedro, J.G. & Barbosa, L.A.G. (1955). A Vegetação. Esboço de Reconhecimento Ecológica-Agrícola de Moçambique. Centro de Investigação Científica Algodoeira, Lourenço Marques.
- Spottiswoode, C.N., Patel, I.H., Hermann, E., Timberlake, J.R. & Bayliss, J. (2006). Threatened bird species on two little-known mountains (Mabu and Chipirone) in northern Mozambique. Paper submitted to Ostrich for publication.
- Stattersfield, A.J., Crosby, M.J., Long A.J. & Wege, D.C. (1998). *Endemic Bird Areas of the World: Priorities for biodiversity conservation*, BirdLife International, Cambridge.
- Strugnell, A.M. (2006). A checklist of the spermatophytes of Mount Mulanje, Malawi. *Scripta Botanica Belgica* 34. National Botanic Garden, Meise, Belgium.
- Vincent, J. (1933). The birds of Northern Portuguese East Africa. Comprising a list of, and observations on, the collections made during the British Museum Expedition of 1931–32. Part 1. *Ibis* **13**: 611–652.
- White, F. (1983). *The Vegetation of Africa*. Natural Resources Research No.20. UNESCO, Paris.
- White, F., Dowsett-Lemaire, F. & Chapman, J.D. (2001). *Evergreen Forest Flora of Malawi*. Royal Botanic Gardens Kew, London.
- Wild, H. & Barbosa, L.A.G. (1968). Vegetation map of the Flora Zambesiaca area., M.O. Collins, Harare.

Suggested citation: Timberlake, J.R., Bayliss, J., Alves T., Baena, S., Francisco, J., Harris, T. & da Sousa, C. (2007). The Biodiversity and Conservation of Mount Chipirone, Mozambique. Report produced under the Darwin Initiative Award 15/036. Royal Botanic Gardens, Kew, London. pp. 33.

## ANNEX 1. Participants on Mt Chipirone expedition, Nov/Dec 2006

Tereza Alves, Forestry Research Section, IIAM, Marracuene, Mozambique.

Aurélio Constantino Banze, Plant Collector, National Herbarium, IIAM, Maputo, Mozambique.

Julian Bayliss, Darwin Project Regional Coordinator, Mulanje Mountain Conservation Trust, Mulanje, Malawi.

Domingos de Azevedo Chiconca, Pastures Section, Chobela Livestock Research Station, IIAM, Chobela, Mozambique.

Jorge R. Francisco, GIS Unit, Land & Water Department, IIAM, Maputo, Mozambique.

Tim Harris, Herbarium, RBG Kew, London, UK.

Rogério Mauro da Costa Jamice, Station Manager, Madonge Forest Research Station, IIAM, Chimoio, Mozambique.

Ivete Frederico Maluleque, Forestry Technician, Regional Research Station, IIAM, Nampula, Mozambique.

Celestino Moises, Driver, IIAM, Chimoio, Mozambique.

Steven Mphamba, Herbarium Assistant, Forestry Research Institute of Malawi, Zomba, Malawi.

David Nangoma, Ecologist, Mulanje Mountain Conservation Trust, Mulanje, Malawi. Email:

Artur Olisse, Driver, IIAM, Maputo, Mozambique.

Hassam Patel, Botanist, Mulanje Mountain Conservation Trust, Mulanje, Malawi.

Camila de Sousa, Forestry Research Section, IIAM, Marracuene, Mozambique.

Christopher Tembo, Herbarium Assistant, National Herbarium & Botanic Gardens, Zomba, Malawi.

Jonathan Timberlake, Project Coordinator, Herbarium, RBG Kew, London, UK.

## ANNEX 2. Plant checklist from Mt Chipirone above 800 m.

T - tree; S - shrub; t - small tree; cl - liana; h - herb; gr - grass; ps - parasite; ge - geophyte;  
su - succulent; ep – epiphytic

MF-h - moist forest, high altitude; MF-m - moist forest, medium altitude; W - miombo woodland;

D/F - heavily disturbed/fallow; Shrub - shrubland; R - riverine

Family	Species	Alt.	L/F	Habitat	Conf.	Notes
<b>PTERIDOPHYTA</b>						
Fern	<i>Asplenium anisophyllum Kunze</i>	1000-1800	h	MF-h	✓	
Fern	<i>Asplenium dregeanum Kunze</i>	1000	h	MF-m	✓	
Fern	<i>Asplenium sandersonii Hook.</i>	1800	h	MF-m/h	✓	
Fern	<i>Marattia fraxinea Sm.</i>	1000	h	MF-m	✓	
Fern	<i>Mohria lepigera (Baker) Baker</i>	2000	h	Shrub	✓	
Fern	<i>Oleandra distenta Kunze</i>	1800	ep	W	✓	
Fern	<i>Pellaea doniana Hook.</i>	1000	h	W	✓	
Fern	<i>Polypodium polypodioides (L.) Hitchcock subsp. ecklonii (Kunze) Schelpe</i>	1000	ep	W	✓	
Pteridaceae	<i>Pteridium aquilinum (L.) Kuhn</i>	1000	h	W	✓	
Fern	<i>Selaginella kraussiana (Kunze) A.Braun</i>	1800	h	MF-h	✓	
<b>MONOCOTYLEDONS</b>						
Aloaceae	<i>Aloe arborescens Mill.</i>	2000	su	Shrub	✓	summit
Aloaceae	<i>Aloe sp.</i>	1000	su	W		miombo
Amaryllidaceae	<i>Scadoxus multiflorus (Martyn) Raf. subsp. multiflorus</i>	1000	ge	D/F	✓	
Araceae	<i>Culcasia falcifolia Engl.</i>	1000	cl	MF-m	✓	
Araceae	<i>Gonatopus clavatus Mayo</i>	at 500m	h	D/F	✓	
Asparagaceae	<i>Asparagus africanus Lam. var. africanus</i>	1000	h	W	✓	
Asparagaceae	<i>Asparagus virgatus Baker</i>	800	h	D/F	✓	
Behniaceae	<i>Behnia reticulata (Thunb.) Didr.</i>	1000	cl	MF-m	✓	
Commelinaceae	<i>Commelina zambesica C.B. Clarke</i>	1000	h	MF-m	✓	
Commelinaceae	<i>Murdannia simplex (Vahl) Brenan</i>	1000	h	W	✓	
Commelinaceae	<i>Pollia condensata C.B. Clarke</i>	1000	h	MF-m	✓	new to FZ area
Cyperaceae	<i>Cyperus cf. amauropus Steud.</i>	1000	h	W	✓	new to Moz
Cyperaceae	<i>Cyperus hemisphaericus Boeck.</i>	1000	h	W	✓	
Cyperaceae	<i>Cyperus pseudoleptocladus Kule</i>	1800	h	W	✓	
Cyperaceae	<i>Cyperus sp.</i>	1000	h	W		
Cyperaceae	<i>Coleochloa cf. setigera (Ridl.) Gilly</i>	2000	h	Shrub	✓	
Dracaenaceae	<i>Dracaena fragrans (L.) Ker Gawl.</i>	1000	S	MF-m	✓	new to Moz Z:
Dracaenaceae	<i>Dracaena laxissima Engl.</i>	1800	S	MF-h	✓	
Dracaenaceae	<i>Dracaena mannii Baker</i>	1000	S	MF-m	✓	
Iridaceae	<i>Gladiolus sp.</i>		ge	W		
Hyacinthaceae	<i>Drimia altissima (L.f.) Ker Gawl.</i>	800	ge	W	✓	
Hyacinthaceae	<i>Ledebouria revoluta (L.f.) Jessop</i>	1000	ge	D/F	✓	
Hypoxidaceae	<i>Hypoxis angustifolia Lam.</i>	1000	h	W	✓	
Musaceae	<i>Ensete ventricosum (Welw.) Cheesman</i>	1000	S	MF-m, W	✓	
Orchidaceae	<i>Aerangis sp.</i>	1000	ep	W		
Poaceae	<i>Andropogon schirensis Hochst.</i>	1000	gr	W	✓	
Poaceae	<i>Hyparrhenia sp.</i>	1000	gr	D/F		
Poaceae	<i>Leptaspis cochleata Thwaites</i>	1000	gr	MF-m	✓	
Poaceae	<i>Oxytenanthera abyssinica (A.Rich.) Munro</i>	1000	S	D/F	✓	
Poaceae	<i>Pennisetum purpureum Schumach.</i>	1000	gr	D/F	✓	
Smilacaceae	<i>Smilax anceps Willd.</i>	1000	cl	D/F	✓	
Zingiberaceae	<i>Afromomum albiflorum Lock</i>	1000	ge	W	✓	
Zingiberaceae	<i>Siphonochilus aethiopicus (Schweinf.) B.L.Burt</i>	1000	ge	W	✓	
Zingiberaceae	<i>Siphonochilus kirkii (Hook.f.) B.L.Burt</i>	1000	ge	W	✓	
<b>DICOTYLEDONS</b>						
Acanthaceae	<i>Crossandra pyrophila Vollesen</i>	1000	h	D/F	✓	
Acanthaceae	<i>Pseuderanthemum subviscosum (C.B. Clarke) Stapf</i>	1000	S	MF-m	✓	
Acanthaceae	<i>Ruellia prostrata Poir.</i>	850	h	W	✓	
Acanthaceae	<i>Thunbergia lancifolia T.Anders.</i>	850	h	W	✓	
Amaranthaceae	<i>Achyranthes aspera L. var. pubescens (Moq.) Townsend</i>	1000	h	W	✓	
Amaranthaceae	<i>Achyranthes aspera L. var. sicula L.</i>	1000	h	D/F	✓	
Anacardiaceae	<i>Lannea discolor (Sond.) Engl.</i>	1000	T	W	✓	

Family	Species	Alt.	L/F	Habitat	Conf.	Notes
Anacardiaceae	<i>Ozoroa insignis Delile</i> subsp. <i>reticulata</i> (Baker f.) Gillett	at 500m	t	D/F	✓	
Annonaceae	<i>Annona senegalensis Pers.</i>	1000	t	D/F	✓	
Apiaceae	<i>Heteromorpha arborescens (Spreng.) Cham. &amp; Schltd.</i> var. <i>montana P.J.D. Winter</i>	1000	t	W	✓	
Apiaceae	<i>Steganotaenia araliacea Hochst.</i>	1000	t	W, D/F	✓	
Apocynaceae	<i>Carissa bispinosa (L.) Brenan</i> subsp. <i>zambesiensis Kupicha</i>	1500	S	W	✓	
Apocynaceae	<i>Carvalhoa campanulata K.Schum.</i>	1000	S	MF-m	✓	
Apocynaceae	<i>Diplorhynchus condylocarpon (Müll.Arg.) Pichon</i>	1000	T	W	✓	
Apocynaceae	<i>Funtumia africana (Benth.) Stapf</i>	1000	T	MF-m	✓	
Apocynaceae	<i>Landolphia sp.</i>		cl	MF-m		
Apocynaceae	<i>Pleiocarpa pycnantha (K.Schum.) Stapf</i>	1000	T/S	MF-m	✓	
Apocynaceae	<i>Rauvolfia caffra Sond.</i>		T	MF-m	✓	
Apocynaceae	<i>Tabernaemontana stapfiana Britten</i>	1800	T	MF-h	✓	
Araliaceae	<i>Cussonia arborea A.Rich.</i>	1000	T	W, D/F	✓	
Araliaceae	<i>Cussonia spicata Thunb.</i>	1000	T	W/F	✓	
Araliaceae	<i>Polyscias fulva (Hiern) Harms</i>	1000	T	MF-m	✓	
Araliaceae	<i>Schefflera goetzenii Harms</i>	1800	T	MF-h,W	✓	
Asclepiadaceae	<i>Glossostelma carsonii (N.E.Br.) Bullock</i>	1000	h	W	✓	
Asclepiadaceae	<i>Margaretta rosea Oliv.</i> subsp. <i>whytei (K.Schum.) Mwanyambo</i>	700	h	D/F	✓	
Asteraceae	<i>Berkhaya zeyheri Oliv. &amp; Hiern</i>	1000	h	W	✓	
Asteraceae	<i>Conyza bonariensis (L.) Cronquist</i>	1000	h	D/F	✓	
Asteraceae	<i>Gerbera viridifolia (DC.) Sch.</i>	1000	h	W	✓	
Asteraceae	<i>Nidorella auriculata DC.</i>	1000	h	D/F	✓	
Balanophoraceae	<i>Sarcophyte sanguinea Sparrm.</i> subsp. <i>sanguinea</i>	1000	ps	W	✓	
Balsamaceae	<i>Impatiens walleriana Hook.f.</i>	1000-1500	h	MF-m	✓	
Bignoniaceae	<i>Markhamia obtusifolia (Baker) Sprague</i>	1000	t	W	✓	
Campanulaceae	<i>Wahlenbergia abyssinica (A.Rich.) Thulin</i> subsp. <i>abyssinica</i>	1000	h	D/F	✓	
Cecropiaceae	<i>Myrianthus holstii Engl.</i>	1000-1500	T	MF-m	✓	
Celastraceae	<i>Catha edulis (Vahl) Endl.</i>	1000	T	MF-m,W	✓	
Celastraceae	<i>Maytenus acuminata (L.f.) Loes.</i> var. <i>acuminata</i>	1800	t	MF-h	✓	
Celastraceae	<i>Maytenus undata (Thunb.) Blakelock</i>	1800	t	MF-h	✓	
Chrysobalanaceae	<i>Parinari excelsa Sabine</i>	1000	T	W	✓	
Clusiaceae	<i>Garcinia kingaensis Engl.</i>	1000-1500	T	MF-m/h	✓	
Clusiaceae	<i>Garcinia volkensii Engl.</i>	1000	T	MF-m	✓	
Clusiaceae	<i>Harungana madagascariensis Poir.</i>	1000-1600	t	MF-m	✓	
Clusiaceae	<i>Psorospermum febrifugum Spach</i>	1000	t	W	✓	
Combretaceae	<i>Combretum molle G.Don.</i>	1000	t	W, D/F	✓	
Connaraceae	<i>Agelaea pentagyna (Lam.) Baill.</i>	1500	cl	MF-m	✓	
Convolvulaceae	<i>Astripomoea malvacea (Klotzsch) Meeuse</i> var. <i>malvacea</i>	1000	h	W	✓	
Crassulaceae	<i>Crassula swaziensis Schonl.</i>	2000	su	Shrub	✓	
Cucurbitaceae	<i>Coccinea adoensis Cogn.</i>	800	h	W	✓	
Cucurbitaceae	<i>Momordica foetida Schumach.</i>	1000	h	D/F	✓	
Dipterocarpaceae	<i>Monotes africanus A.DC.</i>	1000	T	W	✓	
Ebenaceae	<i>Diospyros abyssinica (Hiern) F.White</i> subsp. <i>attenuata F.White</i>	1000-1800	T	MF-m	✓	
Ebenaceae	<i>Diospyros whyteana (Hiern) F.White</i>	2000	t	MF-h	✓	
Ericaceae	<i>Erica cf. johnstoniana Britten</i>	2000	S	Shrub	✓	
Erythroxylaceae	<i>Erythroxylum emarginatum Thonn.</i>	1000-2000	t	W	✓	
Euphorbiaceae	<i>Acalypha villicaulis Hochst.</i>	1000	h	W	✓	
Euphorbiaceae	<i>Acalypha welwitschiana Müll.</i>	1800	S	W	✓	
Euphorbiaceae	<i>Antidesma membranaceum Müll.</i>	1000	t	W	✓	
Euphorbiaceae	<i>Bridelia micrantha (Hochst.) Baill.</i>	1000	T	W,D/F	✓	
Euphorbiaceae	<i>Croton sylvaticus Hochst.</i>	1800	t	MF-h	✓	
Euphorbiaceae	<i>Drypetes gerrardii Hutch.</i> var. <i>grandifolia Radcl.-Sm.</i>	1000-1800	T	MF-mh	✓	
Euphorbiaceae	<i>Drypetes sp.</i>	1000	T	MF-m		unmatched
Euphorbiaceae	<i>Erythrocoeca polyandra (Pax &amp; K.Hoffm.) Prain</i>	1000	t	W, MF-m	✓	
Euphorbiaceae	<i>Macaranga capensis (Baill.) Sim</i>	1000	T	MF-m	✓	
Euphorbiaceae	<i>Margaritarea discoidea (Baill.) G.L.Webster</i> var. <i>nitida (Pax) Radcl.-Sm.</i>	600	t	D/F	✓	
Euphorbiaceae	<i>Neoboutonia macrocalyx Pax</i>	1000	t	MF-m	✓	
Euphorbiaceae	<i>Uapaca kirkiana Müll.Arg.</i>	1000	T	W	✓	
Euphorbiaceae	<i>Uapaca nitida Müll.Arg.</i> var. <i>nitida</i>	1000	t	W	✓	
Fab: Caesalpinioideae	<i>Brachystegia boehmii Taub.</i>	at 500m	T	W	✓	
Fab: Caesalpinioideae	<i>Brachystegia spiciformis Benth.</i>	1000	T	W	✓	
Fab: Caesalpinioideae	<i>Brachystegia tamarindoides Benth.</i> subsp. <i>microphylla (Harms) Chikuni</i>	1000	T	W	✓	

Family	Species	Alt.	L/F	Habitat	Conf.	Notes
Fab: Caesalpinioideae	<i>Brachystegia utilis</i> Hutch. & Burt Davy	1000	T	W	√	
Fab: Caesalpinioideae	<i>Julbernardia globiflora</i> (Benth.) Troupin	1000	T	W	√	
Fab: Mimosoideae	<i>Acacia abyssinica</i> Benth.	1200	T	W	√	
Fab: Mimosoideae	<i>Acacia nilotica</i> (L.) Delile subsp. <i>kraussiana</i> (Vatke) Brenan	at 500m	T	D/F	√	
Fab: Mimosoideae	<i>Albizia adianthifolia</i> (Schumach.) W. Wight	1000	T	MF-m	√	
Fab: Mimosoideae	<i>Dichrostachys cinerea</i> (L.) Wight & Arn. subsp. <i>africana</i> Brenan & Brummitt	500m	S	W	√	
Fab: Mimosoideae	<i>Elephantorrhiza goetzei</i> (Harms) Harms subsp. <i>goetzei</i>	at 500m	S	D/F	√	
Fab: Mimosoideae	<i>Entada rheedei</i> Spreng.	800	cl	W	√	
Fab: Mimosoideae	<i>Newtonia buchananii</i> (Baker) G.C.C. Gilbert & Boutique	1000	T	MF-m	√	
Fab: Papilionioideae	<i>Abrus melanospermus</i> Hassk. subsp. <i>suffruticosus</i> (Boutique) D.K. Harder	1000	h	W	√	new to Moz
Fab: Papilionioideae	<i>Aeschynomene nyassana</i> Taub.	1000	h	W	√	
Fab: Papilionioideae	<i>Dalbergia nitidula</i> Baker	at 500m	t	W, D/F	√	
Fab: Papilionioideae	<i>Desmodium gangeticum</i> (L.) DC.	800	h	W	√	
Fab: Papilionioideae	<i>Dolichos kilimandscharicus</i> Taub. subsp. <i>kilimandscharicus</i>	800	h	D/F	√	
Fab: Papilionioideae	<i>Erythrina abyssinica</i> DC.	1000	T	W	√	
Fab: Papilionioideae	<i>Millettia stuhlmannii</i> Taub.	at 500m	T	W	√	
Fab: Papilionioideae	<i>Mucuna pruriens</i> (L.) DC. var. <i>pruriens</i>	1000	cl	D/F	√	
Fab: Papilionioideae	<i>Pericopsis angolensis</i> (Baker) Meeuwen	1000	T	W	√	
Fab: Papilionioideae	<i>Pterocarpus angolensis</i> DC.	1000	T	W	√	
Flacourtiaceae	<i>Dovyalis macrocalyx</i> (Oliv.) Warb.	1000-1800	S	MF-h, W	√	
Flacourtiaceae	<i>Flacourtia indica</i> (Burm.f.) Merr.	600	t	W/D	√	
Flacourtiaceae	<i>Rawsonia lucida</i> Harv. & Sond.	1000-1500	t	MF-mh	√	
Gesneriaceae	<i>Streptocarpus</i> cf. <i>goetzei</i> Engl.	1500	ep	MF-h	√	
Lamiaceae	<i>Achryospermum laterale</i> Baker	1500	h	W	√	
Lamiaceae	<i>Leucas milanjiana</i> Gürke	1000	h	W, D/F	√	
Lamiaceae	<i>Ocimum obovatum</i> Benth. subsp. <i>obovatum</i>	1000	h	W	√	
Lamiaceae	<i>Plectranthus</i> cf. <i>hadiensis</i> (Forssk.) E.A. Bruce	1000	h	W	√	
Lamiaceae	<i>Plectranthus kapatensis</i> (R.E. Fr.) J.K. Morton	1000	h	W	√	new to Moz Z:
Lamiaceae	<i>Scutellaria schweinfurthii</i> Briq. subsp. <i>paucifolia</i> (Baker) Paton	1000	h	MF-m	√	
Lamiaceae	<i>Vitex</i> cf. <i>doniana</i> Sweet	1000	t	W	√	
Loganiaceae	<i>Anthocleista grandiflora</i> Gilg	1000	T	MF-m	√	
Loganiaceae	<i>Buddleja salviifolia</i> (L.) Lam.	1800	t	Shrub	√	
Loganiaceae	<i>Mostuea brunonis</i> Didr. var. <i>brunonis</i>	1800	S	MF-h	√	new to Moz Z:
Loganiaceae	<i>Nuxia floribunda</i> Benth.	1500	t	MF-m	√	
Loganiaceae	<i>Strychnos spinosa</i> Lam.	1000	t	W	√	
Loganiaceae	<i>Strychnos</i> sp. - forest climber	1000	cl	MF-m		unmatched
Loranthaceae	<i>Agelanthus subulatus</i> (Engl.) Polhill & Wiens	1000	ep	W	√	
Malvaceae	<i>Hibiscus fuscus</i> Garcke	1000	h	D/F	√	
Melastomataceae	<i>Dissotis</i> sp.	1000	S	D/F		
Melastomataceae	<i>Dissotis</i> sp.	2000	S	Shrub	√	
Meliaceae	<i>Ekebergia capensis</i> Sparm.	1800	T	MF-h	√	
Meliaceae	<i>Khaya anthotheca</i> (Welw.) C. DC.	1000-1500	T	MF-m	√	
Meliaceae	<i>Trichilia dregeana</i> Sond.	1000	T	MF-m	√	
Monimiaceae	<i>Xymalos monospora</i> (Harv.) Warb.	1800	T	MF-h	√	
Moraceae	<i>Ficus exasperata</i> Vahl.	1000	T	W	√	
Moraceae	<i>Ficus scassellatii</i> Pamp.	1000	T	MF-m	√	
Moraceae	<i>Ficus</i> sp. - strangling	1000	T	MF-m		
Moraceae	<i>Ficus</i> sur Forssk.	1000	T	W	√	
Moraceae	<i>Treulia africana</i> Decne subsp. <i>africana</i> var. <i>africana</i>	at 500 m	T	W	√	
Moraceae	<i>Trilepisium madagascariense</i> DC.	1000	T	MF-m	√	
Myricaceae	<i>Morella pilulifera</i> (Rendle) Killick	1000	t	W	√	
Myrsinaceae	<i>Maesa lanceolata</i> Forssk.	1000	t	W	√	
Myrsinaceae	<i>Myrsine africana</i> L.	2000	S	MF-h	√	
Myrsinaceae	<i>Rapanea melanophloeos</i> (L.) Mez	1000	T	W	√	
Myrtaceae	<i>Psidium guajava</i> L.	1000	t	D/F	√	
Myrtaceae	<i>Syzygium cordatum</i> C. Krauss	1000	T	W	√	
Myrtaceae	<i>Syzygium guineense</i> (Willd.) DC. subsp. <i>afromontanum</i> F. White	1000-1800	T	MF-mh	√	
Ochnaceae	<i>Ochna holstii</i> Engl.	1800	t	MF-h	√	
Ochnaceae	<i>Ochna</i> cf. <i>mossambicensis</i> Klotzsch	at 500m	S	D/F	√	
Olacaceae	<i>Strombosia scheffleri</i> Engl.	1000-1600	T	MF-m	√	
Olacaceae	<i>Ximenia caffra</i> Sond. var. <i>natalensis</i> Sond.	at 500m	S	W	√	
Oleaceae	<i>Olea capensis</i> L. subsp. <i>macrocarpa</i> (C.H. Wright) I. Verd.	1600	T	MF-h	√	
Oleaceae	<i>Schrebera trichoclada</i> Welw.	at 500 m	t	D/F	√	

Family	Species	Alt.	L/F	Habitat	Conf.	Notes
Passifloraceae	<i>Adenia digitata</i> (Harv.) Engl.	1000	h	W	✓	
Passifloraceae	<i>Adenia rumicifolia</i> Engl. & Harms var. <i>rumicifolia</i>	1000	h	MF-m	✓	
Piperaceae	<i>Piper capensis</i> L.f.	1000	S	MF-m	✓	
Polygalaceae	<i>Securidaca longepedunculata</i> Fresen.	at 500m	T	W	✓	
Proteaceae	<i>Faurea saligna</i> Harv.	1000	T	W	✓	
Proteaceae	<i>Protea welwitschii</i> Engl.	1000	t	W	✓	
Rubiaceae	<i>Chassalia parvifolia</i> K.Schum.	1000	S	MF-m	✓	
Rubiaceae	<i>Coffea mufindiensis</i> Bridson subsp. <i>australis</i> Bridson	1000-1500	S	MF-m	✓	VU
Rubiaceae	<i>Coffea salvatrix</i> Swynn. & Phillipson	1000	t	MF-m	✓	
Rubiaceae	<i>Cremospora triflora</i> (Thonn.) K.Schum.	1000	S	MF-m	✓	
Rubiaceae	<i>Lasianthus kilimandscharicus</i> K.Schum.	2000	S	MF-h	✓	
Rubiaceae	<i>Mussaenda arcuata</i> Poir.	1000	S	W	✓	
Rubiaceae	<i>Oleandra distenta</i> Kunze	1800	S	W	✓	
Rubiaceae	<i>Oxyanthus speciosus</i> DC. subsp. <i>stenocarpus</i> (K.Schum.) Bridson	1000	S	MF-m	✓	
Rubiaceae	<i>Pauridiantha symplocoides</i> (S.Moore) Bremek.	1800	S	W	✓	
Rubiaceae	<i>Pavetta chapmannii</i> Bridson	1800	S	W	✓	
Rubiaceae	<i>Pavetta</i> sp. A	1600	S	W	✓	
Rubiaceae	<i>Psychotria zombamontana</i> (Kuntze) Petit	1600	t	MF-h,W	✓	
Rubiaceae	<i>Pyrostria</i> sp.	1800	S	MF-h	✓	
Rubiaceae	<i>Rothmannia urcelliformis</i> (Hiern) Robyns	1000	T	MF-m	✓	
Rubiaceae	<i>Rytigynia uhligii</i> (K.Schum. & K.Krause) Verdc.	1000	t	MF-m	✓	
Rubiaceae	<i>Tricalysia acantheroides</i> K.Schum.	1800	t	W	✓	
Rubiaceae	<i>Rytigynia uhligii</i> (K.Schum. & K.Krause) Verdc.	1000	S	MF-m	✓	
Rubiaceae	<i>Vangueria infausta</i> Burchell subsp. <i>rotundata</i> (Robyns) Verdc.	1000	t	W	✓	
Rutaceae	<i>Clausena anisata</i> (Willd.) Benth.		S	W		
Rutaceae	<i>Toddalia asiatica</i> (L.) Lam.	1000	cl	MF-m	✓	
Rutaceae	<i>Vepris</i> cf. <i>bachmannii</i> (Engl.) W.Mziray	1500-2000	t	MF-h	✓	
Rutaceae	<i>Vepris nobilis</i> (Delile) W.Mziray	1600	t	MF-h	✓	
Rutaceae	<i>Zanthoxylum gilletti</i> (De Wild.) P.G.Waterman	1000	T	MF-m		
Sapindaceae	<i>Allophylus</i> cf. <i>chaunostachys</i> Gilg	1500-2000	S	MF-h	✓	
Sapindaceae	<i>Blighia unijugata</i> Baker	1000	t	MF-m	✓	
Sapindaceae	<i>Dodonea viscosa</i> Jacq.	2000	S	Shrub	✓	uncertain
Sapindaceae	<i>Macphersonia gracilis</i> O.Hoffm. var. <i>hildebrandtii</i> (O.Hoffm.) Capuron	1500	S	MF-h	✓	
Sapotaceae	<i>Chrysophyllum gorungosanum</i> Engl.	1000	T	MF-m	✓	
Sapotaceae	<i>Englerophytum magalimontanum</i> (Sond.) T.D.Penn.	1000	T	W	✓	
Sapotaceae	<i>Englerophytum natalense</i> (Sond.) T.D.Penn.	1000	T	MF-m	✓	
Sapotaceae	<i>Synsepalum brevipes</i> (Baker f.) T.D.Penn.	at 500m	T	R	✓	
Sapotaceae	<i>Synsepalum muelleri</i> (Kupicha) T.D.Penn.	1000-1800	t	MF-mh	✓	
Scrophulariaceae	<i>Cycnium adonense</i> Benth.	1000	h	W	✓	
Simaroubaceae	<i>Harrisonia abyssinica</i> A.Juss.	1000	t	W		
Solanaceae	<i>Solanum</i> cf. <i>giganteum</i> Jacq.	1200-1500	S	MF-mh	✓	
Sterculiaceae	<i>Cola greenwayii</i> Brenan	1000	T	MF-m	✓	
Sterculiaceae	<i>Dombeya burgessiae</i> Harv.	1000	S	D/F	✓	
Thymeleaceae	<i>Peddiea africana</i> Harv.	1500-2000	t	MF-h	✓	
Tiliaceae	<i>Triumfetta</i> cf. <i>pilosa</i> Roth var. <i>effusa</i> (Harv.) Wild	1000	h	D/F	✓	
Turneraceae	<i>Tricliceras longepedunculatum</i> (Mast.) R.Fern.	800	h	D/F	✓	
Ulmaceae	<i>Celtis gomphophylla</i> Baker	1000	T	MF-m	✓	
Ulmaceae	<i>Trema orientalis</i> (L.) Blume	1000	T	D/F	✓	
Violaceae	<i>Rinorea angustifolia</i> (Thouars) Baill. subsp. <i>ardisiiflora</i> (Oliv.) Grey-Wilson	1000-1800	t	MF-m/h	✓	
Violaceae	<i>Rinorea convallarioides</i> (Baker f.) Eyles	1000	T	MF-m	✓	
Violaceae	<i>Rinorea ferruginea</i> Engl.	1000	T	MF-m	✓	
Vitaceae	<i>Cyphostemma</i> cf. <i>congestum</i> (Baker) Descouings	1000	h	D/F	✓	
Vitaceae	<i>Rhoicissus tridentata</i> (L.f.) Wild. & R.B.Drumm.	1000	cl	W	✓	

NB. Altitude given as 1000 m implies can be found from 800–1200 m in suitable habitat.

## ANNEX 3. List of birds seen on Mt Chipirone, 15–18 December 2005 (from Claire Spottiswoode &amp; Eric Herrmann).

Scientific name	Common name
<i>Pernis apivorus</i>	European Honey Buzzard
<i>Buteo augur</i>	Augur Buzzard
<i>Hieraaetus pennatus</i>	Booted Eagle
<i>Francolinus hildebrandti</i>	Hildebrandt's Francolin
<i>Francolinus afer</i>	Red-necked Spurfowl
<i>Treron calvus</i>	African Green Pigeon
<i>Turtur tympanistria</i>	Tambourine Dove
<i>Turtur afer</i>	Blue-spotted Wood Dove
<i>Columba delegorguei</i>	Eastern Bronze-naped Pigeon
<i>Columba larvata</i>	Lemon Dove
<i>Tauraco livingstonii</i>	Livingstone's Turaco
<i>Chrysococcyx klaas</i>	Klaas's Cuckoo
<i>Bubo africanus</i>	Spotted Eagle Owl
<i>Strix woodfordii</i>	African Wood Owl
<i>Macrodipteryx vexillarius</i>	Pennant-winged Nightjar
<i>Apaloderma narina</i>	Narina's Trogon
<i>Bycanistes bucinator</i>	Trumpeter Hornbill
<i>Bycanistes brevis</i>	Silvery-cheeked Hornbill
<i>Stactolaema leucotis</i>	White-eared Barbet
<i>Pogoniulus bilineatus</i>	Yellow-rumped Tinkerbird
<i>Campethera cailliautii</i>	Green-backed Woodpecker
<i>Dendropicos fuscescens</i>	Cardinal Woodpecker
<i>Psalidoprocne orientalist</i>	Eastern Saw-wing
<i>Hirundo abyssinica</i>	Lesser Striped Swallow
<i>Campephaga flava</i>	Black Cuckoo-Shrike
<i>Coracina caesia</i>	Grey Cuckoo-Shrike
<i>Andropadus virens</i>	Little Greenbul
<i>Andropadus importunus</i>	Sombre Greenbul
<i>Phyllastrephus flavostriatus</i>	Yellow-streaked Bulbul
<i>Pycnonotus barbatus</i>	Common Bulbul
<i>Nicator gularis</i>	Eastern Nicator
<i>Cossypha heuglini</i>	White-browed Robin-Chat
<i>Cercomela familiaris</i>	Familiar Chat
<i>Alethe choloensis</i>	Thyolo Alethe
<i>Phylloscopus trochilus</i>	Willow Warbler
<i>Phylloscopus ruficapilla</i>	Yellow-throated Woodland Warbler
<i>Cisticola cantans</i>	Singing Cisticola
<i>Prinia subflava</i>	Tawny-flanked Prinia
<i>Heliolais erythroptera</i>	Red-winged Warbler
<i>Apalis chariessa</i>	White-winged Apalis
<i>Apalis melanocephala</i>	Black-headed Apalis
<i>Camaroptera brachyura</i>	Grey-backed Camaroptera
<i>Bradornis pallidus</i>	Pale Flycatcher
<i>Platysteira peltata</i>	Black-throated Wattle-eye
<i>Turdoides jardineii</i>	Arrow-marked Babbler
<i>Cyanomitra olivacea</i>	Eastern Olive Sunbird
<i>Hedydipna collaris</i>	Collared Sunbird
<i>Cinnyris venusta</i>	Variable Sunbird
<i>Zosterops senegalensis</i>	Yellow White-eye



<i>Telophorus nigrifrons</i>	Black-fronted Bush-Shrike
<i>Tchagra anchietae</i>	Anchieta's Tchagra
<i>Dryoscopus cubla</i>	Black-backed Puffback
<i>Laniarius aethiopicus</i>	Tropical Boubou
<i>Oriolus chlorocephalus</i>	Green-headed Oriole
<i>Dicrurus ludwigii</i>	Square-tailed Drongo
<i>Cinnyricinclus leucogaster</i>	Violet-backed Starling
<i>Ploceus bertrandi</i>	Bertram's Weaver
<i>Ploceus ocularis</i>	Spectacled Weaver
<i>Ploceus bicolor</i>	Dark-backed Weaver
<i>Ploceus olivaceiceps</i>	Olive-headed Weaver
<i>Lagonosticta rhodopareia</i>	Jameson's Firefinch
<i>Estrilda quartinia</i>	Yellow-bellied Waxbill
<i>Estrilda astrild</i>	Common Waxbill
<i>Vidua macroura</i>	Pin-tailed Whydah
<i>Serinus citrinelloides</i>	African Citril
<i>Emberiza cabanisi</i>	Cabanis's Bunting